

Effect of Diet on Depression: A Review of Nutritional Solutions

Hananeh Ahmadnia^{1*}, Hassan Bahrami², Soraya Mohamadzadeh³

1 Department of Biology, Faculty of Basic Sciences, Ferdowsi University of Mashhad, Iran: h.ahmadnia1990@mail.um.ac.ir

2 Independent Health/Nutrition Researcher; hbahrami.research@gmail.com

3 Department of Biology, Faculty of Science, University of Zanjan, Iran, Soraya.mohamadzadeh@znu.ac.ir

*Correspondence to: **Hananeh Ahmadnia** E-mail: h.ahmadnia1990@mail.um.ac.ir

ABSTRACT

Depression is a mental illness with several categories that have common symptoms such as chronic sorrow, lack of interest, lack of pleasure, mood swings between guilt and poor self-esteem, sleep disturbances, and a loss of appetite. Over 300 million individuals globally suffer from depression, and the socioeconomic cost of this debilitating disorder is expected to rise dramatically in the future decades.

Behavioral health illnesses can be prevented and treated to some extent with dietary and nutritional means. Nutritional psychiatry has produced observational and effectiveness evidence regarding the role of healthy dietary patterns in the onset and treatment of depressive symptoms. Diet is associated with depressive symptoms or depression, meaning that an increase in depressive disorders coincide with a deterioration in healthy living choices, including poor quality diets.

Healthy eating habits and adequate intake of essential nutrients via the diet can help prevent and treat depression by reducing symptoms of mental illness. In addition, because nutrition is a modifiable risk factor for depression, it is practical for the public to consider dietary changes to reduce the prevalence of depressive disorders.

This paper reviews the potential value of diet-based actions to manage depression, and ways in which dietary changes could be made to improve mental and cognitive health. Furthermore, some practical solutions for preventing and controlling depression are proposed based on health-related effects of improving dietary habits and life style.

Keywords: Depression, Prevention, Mental Health, Nutrition, Dietary Solutions

INTRODUCTION

Depression is defined as a common mental condition characterized by persistent sadness, a loss of interest and pleasure, variations between feelings of guilt and poor self-esteem, as well as sleep and/or hunger difficulties that interfere with daily functioning. These symptoms appear in many forms of depression and are classified by their strength, duration, and style of presentation (Higuera, 2021).

Depression, anxiety disorders and other mental health conditions are on the rise globally and remain a threat to public health; they are also interlinked to physical health. Some diseases can result in depression, and vice versa. In particular, major depression is a severe, life-threatening, and highly prevalent psychiatric disorder (Bartolomucci & Leopardi, 2009), which may also lead to frequent thoughts about suicide including in young adults (Ljungberg et al., 2020).

Depression may be associated with various risk and protective factors such as biological, psychological, social, environmental, physical activity, smoking, alcohol consumption, sleep quality, and particularly diet. Following the publication of data showing the link between food and mental health, the name "Nutritional Psychiatry" was established to promote and formalize this new field of study (Sarris et al., 2015; Lanuza et al., 2022).

Nutritional psychiatry has produced observational and effectiveness evidence that support the significance of good eating patterns in depression risk and symptom treatment (Marx et al., 2017; Lassale et al., 2019; Firth et al., 2019). The early research in the subject of nutritional psychiatry centered on the function of individual nutrients or single foods in the incidence of mood disorders, as well as the efficacy of separated nutrients (supplements) as therapeutic methods (Marx et al., 2019).

The associations between diet and depressive symptoms are not fully understood due to complications and diversity in the interacting biological processes (Huang et al., 2019). Hence, this review presents some solid evidence pertaining to the link between nutrition and depression, as well as novel nutraceutical treatments and management of those who suffer from such disorders, as a guide for future clinical studies and tailored nutritional interventions.

Nutrients, other dietary substances and depression

Among the various factors associated with depression, understanding the connections between depression and micronutrient deficiencies is significantly important, as deficiency in micronutrients have been related to a variety of health disorders (Wang et al., 2018). The link between depression and macronutrients has been established in several research studies, and the impacts on chronic diseases and weight management have been shown (Larsen et al., 2010; Lagiou et al., 2012). Several reports have indicated a relationship between macronutrient intake and depression, as well as association of diet with prevalence of depression (Oh et al., 2020).

Vitamins

Vitamin B1 (Thiamine) is required for carbohydrate metabolism (Mikkelsen et al., 2016) and it is very important for neuronal cell function because the nervous system is particularly susceptible to oxidative metabolisms and hence to thiamine shortage, as thiamine is the foundation for numerous coenzymes involved in neuronal cell function as well as glucose use. As a result, thiamine deficiency causes a wide range of changes in brain cell functions (Ghaleiha et al., 2016). Neuroinflammation, overexpression of pro-inflammatory cytokines, and arachidonic acid products are all connected to thiamin deficiency, which can cause emotional problems, memory defects, depression, and cognitive decline (Mikkelsen et al., 2017). The link between thiamine shortage and depression, and necessity of sufficient thiamine intake via the diet has been underlined in several studies, as low thiamin levels has commonly been reported in patients with depression

(Mikkelsen, et al., 2016). Some of the best sources of vitamin B1 are peas, beans, whole grain rice, yeast, and nuts, especially raw almonds (Nikseresht et al., 2012). In contrast, conditions that can impair thiamine levels include alcoholism and excessive coffee intake (Bowman, 2018).

The brain needs **niacin** (Vitamin B₃) to get energy and function properly. Niacin deficiency is associated with depression, and numerous research studies have shown that niacin is beneficiary for depression and anxiety symptoms, as well as in individuals with unipolar depressive disorder (Mikkelsen et al., 2016; Prakash et al., 2008; Smesny et al., 2010). An animal model also showed that niacin can help decrease the generation of pro-inflammatory cytokines by macrophages (Lipszyc et al., 2013). As a result, niacin can help decrease inflammation that is a primary cause of depression (Mikkelsen et al., 2017). Some of the best sources of vitamin B₃ are whole wheat, whole grain rice, mushrooms, green peas, potatoes, and avocado. While supplemental doses of the vitamin can have various side effects including liver toxicity, there is no danger in consuming niacin in the amounts found naturally in foods (Julson, 2018).

Vitamin B₆ (Pyridoxine) is important in the regulation of brain functions and mood (Huang et al., 2013). It has an effect on neurotransmitters that govern sadness, pain perception, and anxiety, and deficiencies causes excessive homocysteine levels, which has been related to seizures, migraines, and depression (Hvas et al., 2004). Vitamin B₆ can influence psychotic symptoms in schizophrenia, tiredness, cognitive performance, and depression by lowering homocysteine levels in the blood (Shiloh et al., 2001). Vitamin B₆ is also necessary for the production of GABA, an inhibitory neurotransmitter present in high amounts in brain tissue and the spinal cord (Mikkelsen et al., 2016). Furthermore, plasma pyridoxal phosphate, derived from vitamin B₆, may be substantially lower in patients with depressive symptoms (Hvas et al., 2004). Depressive patients may have higher plasma homocysteine levels associated with poor vitamin B₆ and folate levels (Ebesunun et al., 2012). In contrast, diets rich in Vitamin B₆ and tryptophan can help improve serotonergic neurotransmission in depression caused by a variety of neurodegenerative illnesses (Shabbir et al., 2013). Some of the best sources of vitamin B₃ include carrots, spinach, bananas, green peas, chickpeas, sweet potatoes, and avocado (Whelan, 2017).

Folic acid (folate, vitamin B₉) works closely with vitamins B₆ and B₁₂ to keep metabolic processes running smoothly. Low folate levels are linked to depression (Gilbody, et al., 2007) and can lead to problems with methylation and monoamine metabolism, which can contribute to depression. Folate is required for dopamine production in the brain, and low dopamine levels may worsen depression symptoms (Mikkelsen et al., 2016). Additionally, folate affects noradrenalin and serotonin receptors in the brain, indicating that it has antidepressant properties (Mikkelsen et al., 2017). In patients who have folate deficiency, even antidepressants may not be effective in treatment (Coppen & Bolander-Gouaille, 2005). Folate deficiency has multiple causes, including poor dietary intake, alcoholism, and weakened stomach acid such as due to routine consumption of antacids that affect folate absorption. Some of the best sources of folate are beans, citrus fruits such as lemons, sweet lemons and oranges, green leafy vegetables such as spinach, and nuts such as raw almonds. For most people, it is relatively easy to get enough folic acid through a healthy balanced diet (Cherney, 2019; Watson, 2022).

Vitamin B₁₂ (Cobalamin) is a catch-all term for any molecule with a cobalt-centered corrin nucleus. B₁₂'s function is intertwined with that of folate. Vitamin B₁₂ is essential for the proper

functioning of the brain and neurological system. Vitamin B₁₂ deficiency causes severe depression, suicidal behavior, cognitive decline, irritability, mania, and psychosis. Indeed, up to one-third of depressive individuals have vitamin B₁₂ deficiency, and sufficient levels are linked to improved treatment results (Vogiatzoglou et al., 2008). Fatigue and weakness, constipation, lack of appetite and weight loss, balance difficulties, depression, and cognitive abnormalities are all common clinical signs of vitamin B₁₂ deficiency (Pawlak et al., 2014). Vitamin B₁₂ deficiency has also been linked to bipolar disorder (Permoda-Osip et al., 2013) and this supports the theory that low B₁₂ levels are associated to decreased production of serotonin and other monoamine neurotransmitters (Seppälä et al., 2013). Low vitamin B₁₂ levels are linked to depression, and a deficit, as along with deficiencies of riboflavin and pyridoxine, is associated with major depression. Vitamin B₁₂ and folate are involved in the metabolism of monoamine neurotransmitters such as norepinephrine. Noradrenergic release is inhibited in patients with major depressive disorder (Mikkelsen et al., 2016). Vitamin B₁₂ is not produced by the cells in humans, animals or plants, but by some certain types of bacteria in a healthy gut, which is then absorbed and stored in the body tissues (O'Leary & Samman, 2010). Consuming natural and chemical-free raw vegetables can improve the gut microbiome, providing suitable conditions in which the "good" bacteria can produce significant quantities of vitamin B₁₂ (Watanabe & Bito, 2018). Studies on Iranian villagers with traditional plant-based diets that are very low in animal products have indicated normal vitamin B₁₂ levels (Halsted, 1960). Similarly, vitamin B₁₂ deficiency is relatively uncommon among native Indians where their own local plant-based products are consumed, but the deficiency has been found with significant frequency among vegetarian Indian immigrants in England with similar dietary patterns, possibly because healthy Indian subjects, eating traditional farming products, may have healthier varieties of bacteria in their small intestine than those in the West (Albert, 1980). According to another study in UK, around half of the vegans with entirely plant-based diet, have had sufficient vitamin B₁₂ (Gilsing et al., 2010; Sushma, 2019). This suggests that vitamin B₁₂ deficiency can be prevented by maintaining a healthy gut microbiome. It is noted that taking antacids and overuse of antibiotics may significantly reduce absorption of vitamin B₁₂ (Zeuschner et al., 2013; Sushma, 2019; Ramirez et al., 2020).

Vitamin C (Ascorbic acid) is a water-soluble antioxidant vitamin that is required by the cells in humans and other animals (Lindblad, et al., 2013; Tveden-Nyborg & Lykkesfeldt, 2013). Vitamin C's active form (ascorbate) is known to have free radical scavenging action, interacting with free radicals such as protein thyl radicals. Biomolecules should be protected against oxidative damage by such processes (Nauser et al., 2015). Ascorbic acid also acts as an indirect antioxidant, regenerating vitamin E and assisting in the maintenance of oxidant/antioxidant equilibrium in the plasma membrane. Ascorbic acid's neuroprotective effects have been documented in several studies, and modulation of various neurotransmitter systems, such as dopaminergic, noradrenergic, serotonergic, glutamatergic, and cholinergic systems, has been involved in ascorbic acid's activities in the central nervous system (Moritz et al., 2020). Clinical observations of scurvy patients were used to make the initial connections between ascorbic acid and depression pathogenesis. Notably, depression is known to occur before the physical signs of scurvy, suggesting that vitamin C insufficiency may be frequent among depressed people (Kocot et al., 2017). Furthermore, prolonged deficiency in vitamin C has been linked to decreased activity of the enzyme dopamine β -hydroxylase, which reduces the hydroxylation of dopamine to noradrenaline and, as a result, lowers noradrenaline levels. In rats, ascorbic acid has reduced the comorbidity of type 2 diabetes mellitus and depression (Shivavedi et al., 2019). Also, ascorbic

acid's anti-inflammatory properties may possibly contribute to its antidepressant effect. According to a meta-analysis, serum levels of ascorbic acid in depressed individuals increase following antidepressant treatment, implying that a sufficient intake of vitamin C from natural sources may reverse depression symptoms (Liu et al., 2015). Other research examined the food composition of university students and found a link between depression symptoms and low daily intake of vitamin C, when compared to healthy individuals (Prohan et al., 2014). Sufficient intake of vitamin C rich foods can be recommended as an adjuvant for the treatment of depression based on the data available in the literature (Moritz et al., 2020). Some of the best sources of vitamin C to include in the diet are citrus fruits such as on-tree ripened lemons and oranges, apples, fresh red bell peppers, turnip, and parsley (Kosheleva & Kodentsova, 2013; Chávez et al., 2015; Hill et al., 2018). It is noted that the recommended amount of vitamin C intake for the average person is between 65 to 90 milligrams (mg) a day, and natural food sources easily provide the required levels. But consuming high doses in the form of supplements such as 1000 milligrams of Ascorbic Acid, can lead to development of kidney stones (Thomas et al., 2013), which can be extremely painful and persistently annoying, worsening the severity of depression during the period (Sheng et al., 2017).

Vitamin D (Calciferol), is a fat-soluble steroid hormone necessary for calcium absorption from the stomach into the circulation (Alghamdi et al., 2020). After hydroxylation in the liver, vitamin D is transformed to its active form 25(OH) D3 (1, 25-dihydroxyvitamin D3), commonly known as calcitriol 25(OH)D, which performs a variety of activities, including preserving cellular health by regulating the equilibrium of numerous minerals, such as calcium and phosphate. The National Academy of Medicine considers serum 25(OH)D level in the range of 20-50 ng/ml as sufficient (ideal), 12-20 ng/ml as insufficient, and less than 12 ng/ml as deficient (NIH, 2022). Vitamin D deficiency (but not insufficiency) has been linked to metabolic illnesses such as obesity, diabetes, insulin resistance, and hypertension, as well as neuropsychiatric conditions including schizophrenia, Parkinson's disease, Alzheimer's disease, depression, and cognitive impairment (NIH, 2022; Hussein et al., 2022). Low levels of vitamin D in individuals with recurrent depression imply that they are a high-risk category for vitamin D deficiency that may be a major risk factor for subclinical depression symptoms, according to the findings of a micro-longitudinal observational research among healthy young people (Kaviani et al., 2020). Many tissues, including the brain, generate an active form of vitamin D, as well as vitamin D receptors and vitamin D binding proteins, which have also been found in the central nervous system (CNS), notably in regions linked with mood and depression (Caldwell et al., 2019). Vitamin D as a neurohormone has been linked to neuron cell growth and development, brain function, and mood, as well as neurotransmitter synthesis, release, and regulation (Li et al., 2014; Libuda et al., 2017). Low vitamin D levels are frequent in individuals with depression (Kaviani et al., 2020). Vitamin D metabolites have the ability to cross the blood brain barrier (BBB) (Diesel et al., 2005), and vitamin D receptors are found throughout the brain, including the hippocampus, which has been linked to depression (Menon et al., 2020). As a result, having sufficient vitamin D levels may provide extra therapeutic advantages in the treatment of depression (Sarris et al., 2016), and has shown to help with depression symptoms (Alghamdi et al., 2020), as well as lower incidence of depression (Michele et al., 2020). Most health professionals recommend getting at least 10–30 minutes of sun exposure daily, and/or consuming foods rich in active form of vitamin D. The recommended daily amount for vitamin D is 600 IU (15 mcg). Those living closer to the equator, i.e., in locations that have a latitude less than 37 degrees north of the Earth's equatorial plane, such as the Middle East, North Africa, South Asia, and Central America, may be able to meet their requirements through

sun exposure on clear sunny days of the year. But at high latitudes (such as in Scandinavian countries and Canada), very little vitamin D can be produced by the skin for at least six months of the year, and in case of deficiency, vitamin D supplements may be needed (Leary et al., 2017; Link, 2017; Pratt, 2019; Chauhan et al., 2022). Dietary sources of vitamin D include D₃ in animal products, and D₂ in mushrooms. The naturally-occurring vitamin D from D₃ are found in farm products from the animals that have sufficiently roamed outside in the sunlight, such as eggs from farm chickens, meat from farm lambs, and seafood. Mushrooms also can contain vitamin D, but only after at least 1-2 hours of direct exposure to ultraviolet sunlight in a clear day, but most commercially grown mushrooms are grown in the dark, and thereby, lack vitamin D (Burriesci, 2020; Jones, 2019; Schmid & Walther, 2013).

Vitamin E is an antioxidant vitamin that is lipid-soluble (Miyazawa et al., 2019). Vitamin E possesses antioxidant, anti-inflammatory, and anti-cancer properties in humans, as well as having direct impacts on enzyme activity and gene transcription regulation (Saremi & Arora, 2010). This vitamin is particularly crucial for CNS function, and an increasing number of studies have linked it to neurodegenerative illnesses and mental problems (Muller, 2010). Vitamin E has been shown to modulate synaptic plasticity, particularly adult neurogenesis, which is an important process for cognitive and mood regulation (Ambrogini et al., 2016). Vitamin E has an antidepressant-like impact in depressed animal models by increasing the GSH antioxidant defense system, glutathione peroxidase (GPx), and glutathione reductase (GR) activity in the hippocampus and prefrontal cortex (Farhadnejad et al., 2020). According to some research, there is a link between low vitamin E levels in the blood and severe depression (Desrumaux et al., 2018), and depressed people may have lower vitamin E and vitamin C levels in their blood than healthy people (Farhadnejad et al., 2020). A cross-sectional study of an Iranian population whose average age was 49 years found a link between dietary vitamin E and mild to moderate depression (Banikazemi et al., 2015). Previous cross-sectional studies investigations also found a link between low vitamin E serum/plasma levels and depression in adults, especially in older males aged 60 years or more (Das et al., 2021). Vitamin E may protect against brain damage, and a low dietary intake of the vitamin was linked to mood swings and depression (Banikazemi et al., 2015). According to another study, a lack of antioxidants, notably Zinc and vitamin E, is linked to an increased incidence of depressive symptoms in older males (Das et al., 2021). The amounts of vitamin E and tryptophan were also linked to mental health, implying that sufficient levels of vitamin E and tryptophan, improves mental health. As a result, the authors speculate that the interplay between vitamin E levels and inflammatory pathways may affect the elderly's quality of life (Manosso et al., 2020). The best dietary sources for vitamin E may include natural foods including nuts (such as raw almonds), seeds (particularly sunflower), fruits such as mangos, and vegetables such as turnip (Rizvi et al., 2014). The recommended daily intake of vitamin E for an average-sized adult is 15 mg that can normally be provided by a healthy balanced diet (Manosso et al., 2013).

Minerals

Magnesium ions are the most abundant divalent intracellular cation in human cells, and it plays an important part in a variety of biological activities such as oxidative phosphorylation, energy generation, glycolysis, protein, and nucleic acid synthesis. According to the 2015–2020 Dietary Guidelines for Americans, the optimal Magnesium requirement with food for women is 320 mg/day and 420 mg/day for men (Barbagallo et al., 2021). Magnesium deficiency has been linked to a variety of mental illnesses, including anxiety, depression, and psychotic behavior (Durlach et

al., 2000). In both animal and human studies, a link between magnesium shortage and depression has been shown. Mice fed a Magnesium-deficient diet had behavioral problems that were linked to depression. Similarly, cross-sectional studies have found a negative association between depressed symptoms and Magnesium intake (Wang et al., 2018). In people who suffer from depression, serum Magnesium levels have been found to be lower (Botturi et al., 2020). Magnesium rich foods may be beneficial in the treatment of depression due to association with transduction and biochemical processes linked to depression, as well as the modulatory effect of magnesium on the NMDA receptor complex's ion channel. Sufficient magnesium intakes were linked to less symptoms of depression, according to a comprehensive study (Barbagallo et al., 2021). Some of the best sources of magnesium are dates, raw almonds, whole grain rice, fresh green vegetables (spinach) and seeds such as pumpkin, sesame and sunflowers (Elliott, 2018; Razzaque, 2018).

Zinc is a crucial trace element for numerous biochemical and physiological processes in the brain (Jurowski et al., 2014; Szewczyk et al., 2011), as well as cellular metabolism (Maserejian et al., 2012). Zinc ions modulate numerous ligand- and voltage-gated ion channels in the hippocampus and cortex (Satała et al., 2016), regulating synaptic transmission or acting as neurotransmitters (Wang et al., 2018). Through processes of reduced neurogenesis (Gao et al., 2009; Pfaender et al., 2016) and Neural Plasticity (Pfaender et al., 2016), disruption of Zinc homeostasis in these areas has been linked to a variety of cognitive disturbances, behavioral, and emotional regulation (Takeda, 2000). A certain quantity of zinc is required for the correct maintenance of learning and memory function in the mature brain, and a reduction in synaptic zinc levels can lead to an increase in glutamatergic levels, which are linked to depression (Wang et al., 2018). Zinc deficiency can lower activity of brain derived neurotrophic factor (BDNF), which is one of the most important elements for optimal learning and memory function, which can lead to depression (Młyniec et al., 2015). In both animals and humans, the link between zinc and depression has been explored extensively. Studies on rodents have found a link between zinc deficiency and depressed symptoms. Researchers have also shown that animals who are more resistant to antidepressant therapy had lower serum zinc levels (Wang et al., 2018). According to a meta-analysis of studies, blood zinc concentrations in depressed people were 0.12 µg/mL lower than in healthy people (Swardfager et al., 2013). In experimental rodent studies, when adult rats were fed a zinc-deficient diet, they displayed depression-like behavior (Takeda et al., 2012). In contrast, sufficient zinc consumption was linked to a lower incidence of depressive symptoms, according to cross-sectional research (Szewczyk et al., 2011). Cross-sectional studies have found an inverse relationship between zinc intake and depression symptoms (Szewczyk et al., 2012) and increased depressed symptoms in people with a low zinc diet (Wang et al., 2018). Some of the best sources of zinc are seeds such as pumpkin, sesame and sunflowers, nuts such as raw almonds, vegetables such as green beans and potatoes, whole grain rice, whole grain wheat, lamb, and seafoods such as fish, oysters, shrimp, and mussels (Olza et al., 2017; West, 2018).

Selenium is a trace element that is required for the correct functioning of numerous selenoproteins in the brain and nervous system that are involved in antioxidant defenses. The recommended average daily dose of selenium is 55 µg/day, with optimum serum levels between 70 and 90 µg/L (Wang et al., 2018). Recent research has looked into a link between selenium levels and depression because of its neuromodulatory involvement in brain function (Gao et al., 2012; Pasco et al., 2012). Selenium deficiency was linked to decreasing concentrations of BDNF in rodent research. It is

possible that BDNF concentrations might influence the association between selenium deficiency and depression because it is a neurotrophic factor that has been extensively linked to the pathophysiology of major depressive disorder (Wang et al., 2018). Selenium is required for the correct production and metabolism of thyroid hormones, and thyroid function is linked to neuropsychiatric symptoms such as mood disorders, cognitive impairment, and other psychiatric symptoms (Młyniec et al., 2015). Selenium deficiency and the resulting dysregulation of thyroid function may contribute to the onset of depression. Selenium has been reported to have potent modulatory effects on the dopaminergic, serotonergic, and noradrenergic systems, all of which are implicated in the physiopathology of depression and other mental disorders (Wang et al., 2018). In an intervention trial with mice, treatment with a multi-target selenium-based compound decreased depression symptoms in female mice, indicating that selenium may have an antidepressant effect (Brüning et al., 2011). Poor dietary selenium consumption was linked to a greater risk of developing major depressive disorder, according to data from a case-control study on women aged 20–89 years (Pasco et al., 2012). A cross sectional study that found depressed symptomatology was shown to be enhanced below and above serum selenium levels of 82 and 85 µg/L, with depressive symptomatology being lowest at 85 µg/L (Conner et al., 2015). Some of the best sources of selenium are sunflower seeds, whole grain rice mushrooms, fish, boiled egg, bananas, and spinach (Olsen, 2019).

Iron is a necessary nutrient for humans and serves as a cofactor for numerous enzymes as well as a key component of oxygen transporters in the body (Bao et al., 2012). It has a cofactor function in many processes including the production of tyrosine and tryptophan, which are precursors to the neurotransmitters serotonin, dopamine, and norepinephrine (Richardson et al., 2015). Iron is required for energy generation, DNA synthesis, neurotransmitter synthesis, myelination of neurons (Dusek et al., 2012), and the maintenance of serotonin levels in the brain (Toxqui et al., 2010). Iron and zinc are necessary for cellular function regulation and neuromodulation (Bibi et al., 2020). Many clinical investigations have recently suggested that iron pathology is linked to glutamatergic system dysfunction, as well as malfunctions in the processes and metabolism of cytoarchitectural and glutamate maladaptive alterations in many brain regions that enable cognitive–emotive behaviors (Bibi et al., 2020). In adults, iron deficiency has been linked to depression (Li et al., 2018). In women of reproductive age who are sensitive to depletion of nutritional components, especially iron, a link between inadequate nutrition and depression has been postulated (Bibi et al., 2020). There was a link between serum ferritin, blood hemoglobin, and depression (Pamuk et al., 2016). Iron deficiency anemia also leads to the development of tiredness, anxiety, depression, social, and attention-deficit problems in children (Li et al., 2017). One study indicated that anxiety and depression are more common in children with iron deficiency anemia than in the general population. Anemic patient, particularly one with iron deficiency anemia, exhibits depression-like symptoms in terms of social behavior, gestures, and mood (Noorazar et al., 2015). Overall, it appears that the rise in neurotransmitter concentration is predominantly caused by blood hemoglobin content, with iron-dependent enzymes playing a significant role (Bibi et al., 2020). Not only is iron deficiency potentially connected to the genesis and presentation of depression in the general population, but an iron-rich diet may also help to alleviate depression symptoms (Dama et al., 2018) and lower incidence of depression (Kim et al., 2015; Miki et al., 2015; Li et al., 2018). Some of the best foods rich in iron include date fruits, pistachio, almonds, whole grain bread, red grape, pomegranate, spinach, pumpkin seeds, and red kidney beans (Khan, 2018). Some food sources of iron, also contain some vitamin C that promotes iron absorption, such as raw spinach

that contains 0.8 mg of iron and 8.4 mg vitamin C per cup (Skolmowska et al., 2022; Ware, 2022). Furthermore, adding vitamin C rich sources such as fresh lemon juice to our meal can help improve iron absorption from the foods, as can avoiding coffee and black tea at meal times (Hills, 2019; Fan, 2016; Skolmowska et al., 2022).

Copper is an important trace metal because it is a cofactor for several enzymes involved in biochemical processes including cholesterol, glucose, iron metabolism, cellular respiration, and hormone production (Grubman & White, 2014). Copper concentration and spatial distribution in the brain are unbalanced (Davies et al., 2014) and may fluctuate over time, including during development, aging, and neurodegenerative conditions (Ramos et al., 2014). Copper is essential for the development and operation of the central nervous system and both deficiency and excess can have a significant impact on brain functioning (Scheiber et al., 2014). Copper levels are crucial in oxidative stress processes and have an impact on the catalytic and structural characteristics of several antioxidant enzymes, which might be one of the major causes of depression (Młyniec et al., 2015). The concentration of copper in patients with depression tends to be lower than in those without depression (Ni et al., 2018). Copper consumption was linked to a lower incidence of depression, according to research in Korean teenage females (Kim et al., 2015). The abnormal changes in blood copper levels in individuals with depression, as well as the link between copper and depression etiology, point to its potential as a biomarker and offer a research route for screening and combining useful biomarkers for depression diagnosis and therapy (Ni et al., 2018). Potatoes, nuts, seeds, and whole-grain products are some of the best sources of copper (Pierson et al., 2019).

Amino Acids

N-acetyl cysteine (NAC) lowers inflammatory cytokines, controls glutamate, stimulates neurogenesis, and lowers apoptosis, all of which contribute to depression's neurobiology. As a result, it has numerous possible antidepressant effects (Berk et al., 2013; Michele et al., 2020). Hence, dietary changes towards sufficient intake of proteins, particularly from plant-based sources such as beans and peas, but also from seafood, may help reduce depression (Shen et al., 2021).

S-adenosylmethionine (SAME) is one of the most important molecules that methionine (a sulfur-containing amino acid) can be converted into. In healthy people, SAME is found in safe low levels in almost every tissue and fluid in the body and appears to have a number of potential mechanisms of action, such as improving cell signaling pathways (Lande, 2020; Anjum et al., 2018). However, SAME in higher than normal levels can make it to be a potentially toxic compound, and abnormal levels in the body have been reported in liver diseases and depression (Hopp & Shurtleff, 2017; Tinsley, 2018). SAME is not significantly found in foods, but its precursor, methionine, is plentiful in many protein-rich foods. Although methionine has important roles in the body and is the most variable metabolite found in human plasma, diets that are low in this amino acid are beneficiary for health. Even some cancer cells are dependent on dietary methionine to grow. In these cases, limiting dietary intake of methionine, such as by consuming less animal proteins, could further improve health (Tinsley, 2018; Gao et al., 2019).

Researchers have found that **homocysteine**, an amino acid with significant impact on depression, can be lowered by saffron, therefore, saffron can be considered as a natural antidepressant (Jelodar

et al., 2018). Saffron can be used as a flavoring in food preparations or consumed in the form of tea.

Omega-3 Fatty Acids

Changes in membrane lipids may play a role in the pathophysiology of depression, making lipid-based therapies a viable option (Müller et al., 2015). In the CNS, omega-3 fatty acids have both long-term and short-term effects. Their long-term consequences include an impact on brain structure development, maintenance, and function. The physiology of neuronal signal transduction is affected by omega-3 fatty acids' short-term impacts (Bazinet & Layé, 2014). Animal studies show that omega-3 fatty acids may have an effect on numerous neurobiological mediators thought to be involved in the pathophysiology of MDD (Müller et al., 2015).

Preclinical studies have shown that a diet lacking in omega-3 fatty acids causes depressive-like symptoms and aberrant social behavior in rodents (Bondi et al., 2014). Mice raised on an omega-3 fatty acid-deficient diet, for example, exhibited lower amounts of polyunsaturated fatty acids like docosahexaenoic acid in the brain and higher levels of omega-6 fatty acids, as well as a variety of depressive symptoms in behavioral tests (Lafourcade et al., 2011). Omega-3 fatty acids have also been found to prevent and decrease depression-like behaviors in rat models. However, it's important to remember that most animal models of major depressive illness have significant flaws, especially when it comes to depression subtypes, TRD, and recurrent depression (Lange et al., 2021).

The evidence for using omega-3 fatty acids in the prevention and treatment of depression in humans is based on observational data, including a correlational link between lower prevalence of MDD and higher intake of Omega-3 rich foods such as fish (Ross, 2016). Another study looked at the link between depression and dietary omega-3 fatty acid intake, and found that consumption of Omega-3 rich foods was linked with a substantially lower risk of depression (Grosso et al., 2016). In addition, older individuals in France who eat fish on a regular basis (at least weekly) had less depression symptoms than persons in a control group (Barberger-Gateau et al., 2005). Furthermore, depressed people have lower omega-3 fatty acid levels and a higher omega-6 to omega-3 polyunsaturated fatty acid ratio in their blood and brain (Parletta et al., 2016). A meta-analysis found that omega-3 fatty acids have an overall positive impact on depression symptoms (Liao et al., 2019).

The richest omega-3 sources are certain seafood. Also, regularly consuming sufficient amounts of some plant foods, such as walnut and olive oil that chronic consumption is safe, may provide our daily needs for omega-3 (Wardhana et al., 2011; Sánchez-Villegas et al., 2018; Petrović-Oggiano et al., 2020). In contrast, high dose Omega 3 supplements not only may not be a sustained solution for depression, but they were even found to cause some side effects such as eructation, dyspepsia, diarrhea, vomiting, nausea, and arthralgia, as well as increased risk of aggressive prostate cancer, which all may contribute to worsening depression (Reinberg, 2013; Krupa et al., 2022).

Carotenoids

Carotenoids are a diverse collection of colorful liposoluble pigments found naturally in plants, fungus, bacteria, and algae, as well as in a variety of foods such as fruits, vegetables, and fish

(Milani et al., 2017). 40 carotenoids are present in a typical human diet, but only about around 20 are found in human blood and tissues. Among these carotenoids, β -carotene, α -carotene, lycopene, lutein, and cryptoxanthin are the most significant dietary pigments (Milani et al., 2017). These pigments have been linked to a variety of health benefits, including effects on the neurological system that contribute to neuronal differentiation, neuronal patterning, and motor axon development, due to their powerful anti-inflammatory and antioxidant properties (Ahmed et al., 2021). Inflammation mediates the link between plasma carotenoids and depression. Carotenoid deficiency has been linked to an increased risk of depression, as a result of inflammation or dysregulation of the HPA axis (Milaneschi et al., 2012). Depressed individuals have significantly low levels of antioxidants, especially carotenoids, according to case control studies, and dietary treatment with carotenoids has resulted in a significant decrease in depression (Gautam et al., 2012). Increasing intakes of β -carotene to achieve sufficiency may significantly increase the levels of serotonin and BDNF in individuals suffering from depression (Kim et al., 2015). These findings suggest that sufficient carotenoid levels might help people feel less depressed (Kim et al., 2016) and contribute to treatment of people with depression and anxiety who had low levels of antioxidants in their blood (Gautam et al., 2012). In university students from the United Kingdom, a higher consumption of fruits and vegetables, which are the major source of dietary carotenoids, was found to be inversely associated to various mental health issues (Boozari et al., 2021). Additional data from several groups suggests that carotenoid rich foods is inversely linked with depression risk in both adults and the elderly (Ge et al., 2020; Li & Li, 2019; Milaneschi et al., 2012). Some fruits and vegetables that are rich in dietary carotenoids include red bell peppers, carrots, mangoes, oranges, and spinach (Tan & Norhaizan, 2019).

Polyphenols

Polyphenols are natural chemicals that may be found in a wide range of foods (Pathak et al., 2013). The main groups of polyphenols include phenolic acids, flavonoids, and the less frequent stilbenes and lignans. The majority of polyphenols are digested by colonic bacteria rather than being absorbed via the intestinal barrier (Bayes et al., 2020). Polyphenols preserve mental health by upregulating the body's natural defensive mechanisms, decreasing oxidative damage, and stabilizing free radicals (Gomez-Pinilla & Nguyen, 2012). Polyphenols have been shown in animal tests to decrease depression-like behavior in rodents (Pathak et al., 2013). Polyphenols have also been shown to have neuroprotective effects, regulating particular cellular signaling pathways involved in cognitive functions (Gomez-Pinilla & Nguyen, 2012). Polyphenols' anti-inflammatory characteristics, which include suppression of pro-inflammatory cytokines, free radical scavenging, and antioxidant activity, as well as neuroprotective capabilities, are another proposed explanation for how they benefit mental health (Dias et al., 2012). Metabolic routes and metabolites of polyphenols may potentially be one of the features responsible for their therapeutic benefits, according to research (Bayes et al., 2020). Polyphenols are found in plant-based foods, notably apples, citrus fruit, grapes, whole-grains, nuts and olive oil, and numerous dark colored vegetables, which are common in the Mediterranean diet (Grosso, 2018; Schwingshackl et al., 2020). According to the findings of a meta-analysis, the Mediterranean diet reduces the incidence of depression (Lassale et al., 2019) and improves overall health due to the anti-inflammatory and oxidative stress protective effects of the diet (Lassale et al., 2019; Psaltopoulou et al., 2013).

Probiotics

Probiotics are living microorganisms that colonize the gastrointestinal system, and provide health advantages to the host when consumed in sufficient numbers (Wallace & Milev, 2021; Gawlik-Kotelnicka et al., 2021). Probiotics have been studied in both animal and human research, with promising findings such as reduction of depression- and anxiety-like behavior, meaning that it can act like natural antidepressant in patients with depressed and anxiety symptoms (Gambaro et al., 2020; Wallace & Milev, 2021). Probiotics impacted mice's behavior and reduced symptoms of depression and anxiety (Reininghaus et al., 2020; Peirce & Alviña, 2019). Healthy diets, particularly sufficient intake of fresh vegetables and fruits as a habit, normally can lead to metabolic activity of the human gut microbiota, and natural availability of probiotics in the gut, which in turn can impact health (Conlon et al., 2014; Rinninella et al., 2019). Probiotics can also be acquired via naturally fermented foods such as plain yogurt (Li et al., 2009; Ho & Prasad, 2013).

Dietary components that may worsen depression

In some cross-sectional and prospective studies, dietary patterns such as the Mediterranean diet rich in plant-based foods have been linked to a lower incidence of depression (Lassale et al., 2019). In contrast, eating too few calories, or, excessively consuming animal proteins, may cause metabolic syndrome by slowing down the metabolism, a factor that may be associated with depression (Petre, 2017; Cai et al., 2021).

Consumption of nutrient-rich plant-based foods has significantly decreased in the last few decades because of increased consumption of ultra-processed foods, which involve the removal of key nutrients. Hence it is best for the patients suffering from depression to increase consumption of naturally produced unprocessed foods, in particularly those that can be eaten raw, and minimize consumption of processed fast foods (Razzaque, 2018; Campbell, 2018).

Dietary habits that include plant-based raw foods, such as the traditional African Mediterranean diets, can gradually improve the mood by naturally enhancing the functions related to nervous system and brain, sustainably leading to reduced depression symptoms and anxiety-induced behaviors (Jacka et al., 2017). In addition, if alcohol, addictive substances, refined foods, fast foods, diets containing high levels of trans-fatty acids, sugar-sweetened beverages, and smoking are avoided, depression can be resolved more effectively (Johnson, 2019; Moludi et al., 2020).

Reducing animal protein, and consuming sufficient amounts of fruit, vegetables, and plant-based proteins can help improve mood and reduce depression. Proteins from plants are often lower in methionine than animal proteins, hence plant-based diets could better improve the overall health and increase lifespan (Tinsley, 2018; Gao et al., 2019). In general, diets high in animal proteins may worsen depression, as such proteins are metabolized primarily by microbial fermentation in the intestine, which production of protein fermentation products and metabolites can affect the gut microbiota. The protein fermentation generally forms toxic products, including hydrogen sulfide, amines, and ammonia. Hence, high-protein diets increase the risk of intestinal diseases by producing toxic metabolites in the colon (Cai et al., 2021). In this case, if animal products are consumed excessively, due to increased risk of inflammatory bowel diseases (Wu et al., 2022), the gut microbiota is negatively affected (Alam et al., 2020). The gut microbiota is responsible of regulating availability of tryptophan, and thus the concentration of serotonin (Terry & Margolis,

2016; Carlessi et al., 2021), which have a significant impact on brain serotonin accessibility and depression-like behavior (Lukić et al., 2019; Albenberg & Wu, 2014).

High glycemic-load meals (such as cakes, cookies, sweet treats, fried potatoes, chips) have negative impacts on depression symptoms in non-depressed adults, compared to low glycemic-load meals (such as vegetables, fruits, nuts). Overall, a high glycemic-load diet may cause overall mood changes, increased tiredness, and depressed symptoms (Brey Meyer et al., 2016).

Increased sugar consumption from sugar sweetened drinks and foods has been linked to an increased risk of depression according to prospective cohort studies (Knüppel et al., 2017; Sanchez-Villegas et al., 2018; Guo et al., 2014). A meta-analysis also found a nonlinear dose-response connection between increased sugar-sweetened beverage intake and increased risk of depression (Hu, et al., 2019). A prospective analysis of the Women's Health Initiative cohort found that higher intake of added sugars was linked to an increased risk of incident depression after three years of follow-up (Gangwisch et al., 2015). It should be noted that, as pointed out above, much of the research relating diet to depression could be finding reverse causation. For example, chocolate addiction (which is often consumed with high levels of sugar) is more common in people with untreated chronic depression but this does not inform us which is cause and which is result (Woznicki, 2010).

A diet high in acidogenic (acid-forming) foods (See Bahrami & Greiner, 2021.) may increase the risk of depression, anxiety, stress, and obesity. The researchers found that a high dietary acid load was associated with a higher incidence of psychological disorders. Hence, reducing high dietary acid, such as by sufficient consumption of foods that have alkalizing properties (As explained in Bahrami & Greiner 2021), these are not alkaline foods, can be effective in reducing depression and other mental disorders (Mozaffari et al., 2020). Some foods that form acidic metabolic waste (strong acidifiers) and increase acidity of the blood more significantly include red meats from pork, commercial dairy products, soft drinks, chocolate, coffee, high-fructose corn syrup, table sugar, frying oil, sour plums, peanuts, and fried potatoes (Bahrami and Greiner, 2021; Corriher, 2009). In comparison, plant-based foods result in significantly less acid load (Ronco et al., 2021), and could perhaps be beneficial in treating the depressed patients. Foods that form alkaline metabolic waste include raw almonds, date fruits, olive oil, figs, grapes, lemons, apples, pomegranates, fresh ginger, garlic, onion, carrot, and celery (Bahrami & Greiner, 2021; Bridgeford, 2015). However, bicarbonate supplements and alkaline drinks are not very effective over the long-term (Bahrami & Greiner, 2021).

Hypoxia, poor oxygenation of body tissues including the brain, negatively affects a person's mood and causes depression. Even mild hypoxia may decrease serotonin synthesis, reducing appetite and motivation, and disrupting sleep patterns. Chronic hypoxia in patients with respiratory or cardiac dysfunction could similarly precipitate depression (Kanekar et al., 2015). Healthy dietary habits, as well as aerobic exercise can help improve mood by reducing hypoxia (Kang et al., 2020).

There are now consistent and convincing threads of evidence that support the importance of food and dietary habits as predictors of depression and other prevalent mental maladies. Diet is a very important component of mental health with significant effects on development of depression (Tello, 2020). Some of the diet-related factors that are associated with depression may include

micronutrients deficiency (Ljungberg et al., 2020), metabolic syndrome (Moradi et al., 2021; Cai et al., 2021), and unhealthy diet (Lanuza et al., 2021).

CONCLUSIONS

The existing evidence confirms that nutrient deficiency and unhealthy diets can cause emotional problems, memory defects, depression, and cognitive decline. Healthy dietary patterns can also have a role as an adjunctive treatment for mental disorders; hence, dietary solutions can be considered as novel nutraceutical treatments and management approaches for those who suffer from depression and other mental disorders.

There are a variety of lifestyle and diet modifications that have appropriate potential clinical applications to better manage depression, alongside pharmacotherapies and psychological techniques. Indeed, the time has come for a more integrative approach to treat depression, with acknowledgment of the potential applicability of lifestyle changes and diet modifications. The integrated approach may then allow people live happier, feel less stressed, achieve higher levels of life satisfaction, and face less depression.

In addition to individual solutions, depression should be looked at as a social problem, which requires utilization of healthy diets, not just in individuals, but also in the majority of societies via public training and education, as well as the use of public policy mechanisms such as taxation of harmful foods and subsidization of unprocessed foods.

REFERENCES

- Ahmed, S., Khan, S.T., Aziz, A., Gul, S., Buvnariu, L., Zia-Ul-Haq, M. (2021). 'Role of Carotenoids in Neurological Diseases'. In: Zia-Ul-Haq, M., Dewanjee, S., Riaz, M. (eds) Carotenoids: Structure and Function in the Human Body. Springer, Cham. https://doi.org/10.1007/978-3-030-46459-2_16
- Alam, M. T., Amos, G. C., Murphy, A. R., Murch, S., Wellington, E. M. and Arasaradnam, R. P. (2020) 'Microbial imbalance in inflammatory bowel disease patients at different taxonomic levels', Gut Pathogens, 12(1), pp. 1-8. <https://doi.org/10.1186/s13099-019-0341-6>
- Albenberg, L. G. and Wu, G. D. (2014) 'Diet and the intestinal microbiome: associations, functions, and implications for health and disease', Gastroenterology, 146(6), pp. 1564-1572. <https://doi.org/10.1053/j.gastro.2014.01.058>
- Albert, M., Mathan, V. & Baker, S. 'Vitamin B12 synthesis by human small intestinal bacteria.' Nature 283, pp. 781–782 (1980). <https://doi.org/10.1038/283781a0>
- Alghamdi, S., Alsulami, N., Khoja, S., Alsufiani, H., Tayeb, H. O. and Tarazi, F. I. (2020) 'Vitamin D supplementation ameliorates severity of major depressive disorder', Journal of Molecular Neuroscience, 70(2), pp. 230-235. <https://doi.org/10.1007/s12031-019-01461-2>
- Ambrogini, P., Betti, M., Galati, C., Di Palma, M., Lattanzi, D., Savelli, D., Galli, F., Cuppini, R. and Minelli, A. (2016) 'α-Tocopherol and hippocampal Neural Plasticity in physiological and pathological conditions', International Journal of Molecular Sciences, 17(12), p. 2107. <https://doi.org/10.3390/ijms17122107>
- Anjum, I., Jaffery, S. S., Fayyaz, M., Samoo, Z. and Anjum, S. (2018) 'The role of vitamin D in brain health: a mini literature review', Cureus, 10(7), p. 2960. <https://doi.org/10.7759/cureus.2960>
- Bahrami, H. and Greiner, T. (2021) 'The alkaline diet and the Warburg effect', World Nutrition, 12(1), pp. 20-39. <https://doi.org/10.26596/wn.202112120-39>
- Banikazemi, Z., Mokhber, N., Safarian, M., Mazidi, M., et al. (2015) 'Dietary vitamin E and fat intake are related to Beck's depression score', Clinical Nutrition ESPEN, 10(2), pp. 61-65. <https://doi.org/10.1016/j.clnesp.2014.12.001>
- Bao, W., Rong, Y., Rong, S., and Liu, L. (2012) 'Dietary iron intake, body iron stores, and the risk of type 2 diabetes: a systematic review and meta-analysis', BMC Medicine, 10(1), pp. 1-13. <https://doi.org/10.1186/1741-7015-10-119>
- Barbagallo, M., Veronese, N. and Dominguez, L. J. (2021) 'Magnesium in aging, health and diseases', Nutrients, 13(2), p. 463. <https://doi.org/10.3390/nu13020463>
- Barberger-Gateau, P., Jutand, M. A., Letenneur, L., Larrieu, S., Tavernier, B. and Berr, C. (2005) 'Correlates of regular fish consumption in French elderly community dwellers: data from the Three-City study', European Journal of Clinical Nutrition, 59(7), pp. 817-825. <https://doi.org/10.1038/sj.ejcn.1602145>
- Bartolomucci, A. and Leopardi, R. (2009) 'Stress and depression: preclinical research and clinical implications', PloS one, 4(1), p. e4265. <https://doi.org/10.1371/journal.pone.0004265>

- Bayes, J., Schloss, J. and Sibbritt, D. (2020) 'Effects of polyphenols in a Mediterranean diet on symptoms of depression: a systematic literature review', *Advances in Nutrition*, 11(3), pp. 602-615. <https://doi.org/10.1093/advances/nmz117>
- Bazinet, R. P. and Layé, S. (2014) 'Polyunsaturated fatty acids and their metabolites in brain function and disease', *Nature Reviews Neuroscience*, 15(12), pp. 771-785.
- Berk, M., Malhi, G. S., Gray, L. J. and Dean, O. M. (2013) 'The promise of N-acetylcysteine in neuropsychiatry', *Trends in Pharmacological Sciences*, 34(3), pp. 167-177.
- Bibi, M., Basharat, S., Gilani, S. A., Qamar, M. M., Basharat, A., and Basharat, B. (2020) 'Iron Supplementation Intermittently in Reducing the Severity of Depression', *American Scientific Research Journal for Engineering, Technology, and Sciences*, 69(1), pp. 167-174.
- Bondi, C. O., Taha, A. Y., Tock, J. L., Totah, N. K., Cheon, Y., Torres, G. E., Rapoport, S. I. and Moghaddam, B. (2014) 'Adolescent behavior and dopamine availability are uniquely sensitive to dietary omega-3 fatty acid deficiency', *Biological Psychiatry*, 75(1), pp. 38-46.
- Boozari B, Moradi S, Heydarpour F, Clark CCT, Nezamoleslami S, Saneei P, Safavi SM (2022) The association between carotenoid intake, mental health, and sleep quality among university students. *Sleep Breath*. 2022 Jun;26(2):829-837. <https://doi.org/10.1007/s11325-021-02420-1>
- Botturi, A., Ciappolino, V., Delvecchio, G., Boscutti, A., Viscardi, B. and Brambilla, P. (2020) 'The role and the effect of magnesium in mental disorders: a systematic review', *Nutrients*, 12(6), p. 1661.
- Bowman, J. (2018) What Does Vitamin B-1 Do? <https://www.healthline.com/health/vitamin-watch-b1-thiamine>
- Breymeyer, K. L., Lampe, J. W., McGregor, B. A. and Neuhouser, M. L. (2016) 'Subjective mood and energy levels of healthy weight and overweight/obese healthy adults on high- and low-glycemic load experimental diets', *Appetite*, 107, p. 253-259.
- Bridgeford, R. (2015) 'The definitive acid and alkaline food chart'. <https://liveenergized.com/wp-content/uploads/2018/05/Alkaline-Food-Chart-May.pdf>
- Brüning, C. A., Souza, A. C. G., Gai, B. M., Zeni, G., and Nogueira, C. W. (2011) 'Antidepressant-like effect of m-trifluoromethyl-diphenyl diselenide in the mouse forced swimming test involves opioid and serotonergic systems', *European Journal of Pharmacology*, 658(2-3), pp. 145-149.
- Burriesci H. (2020) How Mushrooms Become High in Vitamin D <https://www.montereymushrooms.com/blog/how-mushrooms-become-high-in-vitamin-d>
- Cai, J., Chen, Z., Wu, W., Lin, Q. and Liang, Y. (2021) 'High animal protein diet and gut microbiota in human health', *Critical Reviews in Food Science and Nutrition*, pp. 1-13.
- Caldwell, J. D., Londe, K., Ochs, S. D., Hajdu, Z., Rodewald, A., Gebhart, V. M., & Jirikowski, G. F. (2019). Three steroid-binding globulins, their localization in the brain and nose, and what they might be doing there. *Steroids*, 142, 48-54.

Campbell, D. (2018) Eating junk food raises risk of depression, says multi-country study. <https://www.theguardian.com/society/2018/sep/26/eating-junk-food-raises-risk-of-depression-says-multi-country-study/>

Carlessi, A. S., Borba, L. A., Zugno, A. I., Quevedo, J. and Réus, G. Z. (2021) 'Gut microbiota–brain axis in depression: The role of neuroinflammation', *European Journal of Neuroscience*, 53(1), pp. 222-235.

Chauhan, K., Shahrokhi, M. and Huecker, M. R. (2022) Vitamin D. Treasure Island (FL): StatPearls Publishing. <https://www.ncbi.nlm.nih.gov/books/NBK441912/>

Chávez-Mendoza, C., Sanchez, E., Muñoz-Marquez, E., Sida-Arreola, J. P. and Flores-Cordova, M. A. (2015) 'Bioactive Compounds and Antioxidant Activity in Different Grafted Varieties of Bell Pepper', *Antioxidants*, 4(2), pp. 427–446.

Cherney, K. (2019) Does Folic Acid Help with Hair Growth <https://www.healthline.com/health/folic-acid-for-hair/>

Conlon MA, Bird AR. (2014) The impact of diet and lifestyle on gut microbiota and human health. *Nutrients*, 7(1):17-44. <https://doi.org/10.3390/nu7010017>

Conner, T. S., Richardson, A. C., and Miller, J. C. (2015) 'Optimal serum selenium concentrations are associated with lower depressive symptoms and negative mood among young adults', *The Journal of Nutrition*, 145(1), pp. 59-65.

Coppen, A. and Bolander-Gouaille, C. (2005) 'Treatment of depression: time to consider folic acid and vitamin B12', *Journal of Psychopharmacology*, 19(1), pp. 59-65.

Corriher, C. S. (2009) PH Food Chart <https://healthwyze.org/reports/191-ph-food-chart/>

Dama, M., Van Lieshout, R. J., Mattina, G. and Steiner, M. (2018) 'Iron deficiency and risk of maternal depression in pregnancy: an observational study', *Journal of Obstetrics and Gynaecology Canada*, 40(6), pp. 698-703.

Das, A., Cumming, R. G., Naganathan, V., Ribeiro, R. V., et al. (2021) 'The association between antioxidant intake, dietary pattern and depressive symptoms in older Australian men: the Concord Health and Ageing in Men Project', *European Journal of Nutrition*, 60(1), pp. 443-454.

Davies, K. M., Bohic, S., Carmona, A., Ortega, R., Cottam, V., Hare, D. J., Finberg, J. P. M., Reyes, S., Halliday, G. M., Mercer, J. F. B., Double, K. L. (2014) 'Copper pathology in vulnerable brain regions in Parkinson's disease', *Neurobiology of Aging*, 35(4), pp. 858-866.

Desrumaux, C. M., Mansuy, M., Lemaire, S., Przybilski, J., Le Guern, N., Givalois, L. and Lagrost, L. (2018) 'Brain vitamin E deficiency during development is associated with increased glutamate levels and anxiety in adult mice', *Frontiers in Behavioral Neuroscience*, 12, p. 310.

Michele, F. D., Talamo, A., Niolu, C. and Siracusano, A. (2020) 'Vitamin D and N-Acetyl Cysteine Supplementation in Treatment-Resistant Depressive Disorder Patients: A General Review', *Current Pharmaceutical Design*, 26, pp. 1-18.

Dias, G.P., Cavegn N, Nix, A., do Nascimento Bevilacqua, M.C., Stangl, D., Zainuddin, M.S., Nardi, A.E., Gardino, P.F., Thuret, S. (2012) 'The role of dietary polyphenols on adult

hippocampal neurogenesis: molecular mechanisms and behavioural effects on depression and anxiety'. *Oxid Med Cell Longevity*, 2012:541971. <https://doi.org/10.1155/2012/541971>

Diesel, B., Radermacher, J., Bureik, M., Bernhardt, R., Seifert, M., Reichrath, J., ... & Meese, E. (2005). 'Vitamin D3 metabolism in human glioblastoma multiforme: functionality of CYP27B1 splice variants, metabolism of calcidiol, and effect of calcitriol'. *Clinical Cancer Research*, 11(15), 5370-5380.

Durlach, J., Bac, P., Bara, M. and Guiet-Bara, A. (2000) 'Physiopathology of symptomatic and latent forms of central nervous hyperexcitability due to magnesium deficiency: a current general scheme', *Magnesium Research*, 13(4), pp. 293-302.

Dusek, P., Jankovic, J. and Le, W. (2012) 'Iron dysregulation in movement disorders', *Neurobiology of Disease*, 46(1), pp. 1-18.

Ebesunun, M. O., Eruvulobi, H. U., Olagunju, T. and Owoeye, O. A. (2012) 'Elevated plasma homocysteine in association with decreased vitamin B12, folate, serotonin, lipids and lipoproteins in depressed patients', *African Journal of Psychiatry*, 15(1), pp. 25-29.

Elliott, B. (2018) '8 Proven Health Benefits of Dates'
<https://www.healthline.com/nutrition/benefits-of-dates/>

Fan, F. S. (2016) 'Iron deficiency anemia due to excessive green tea drinking', *Clinical Case Reports*, 4(11), p. 1053.

Farhadnejad, H., Tehrani, A. N., Salehpour, A. and Hekmatdoost, A. (2020) 'Antioxidant vitamin intakes and risk of depression, anxiety and stress among female adolescents', *Clinical Nutrition ESPEN*, 40, pp. 257-262.

Firth, J., Marx, W., Dash, S., Carney, R., Teasdale, S. B., Solmi, M. Stubbs, B., Schuch, F. B., Carvalho, A. F., Jacka, F., and Sarris, J. (2019) 'The effects of dietary improvement on symptoms of depression and anxiety: a meta-analysis of randomized controlled trials', *Psychosomatic Medicine*, 81(3), p. 265.

Gambaro, E., Gramaglia, C., Baldon, G., Chirico, E., Martelli, M., Renolfi, A. and Zeppego, P. (2020) "'Gut-brain axis": Review of the role of the probiotics in anxiety and depressive disorders', *Brain and Behavior*, 10(10), 01803.

Gangwisch, J. E., Hale, L., Garcia, L., Malaspina, D., Opler, M. G., Payne, M. E., Rossom, R. C. and Lane, D. (2015) 'High glycemic index diet as a risk factor for depression: analyses from the Women's Health Initiative', *The American Journal of Clinical Nutrition*, 102(2), pp. 454-463.

Gao, H. L., Zheng, W., Xin, N., Chi, Z. H., Wang, Z. Y., Chen, J., and Wang, Z. Y. (2009) 'Zinc deficiency reduces neurogenesis accompanied by neuronal apoptosis through caspase-dependent and-independent signaling pathways', *Neurotoxicity Research*, 16(4), pp. 416-425.

Gao, S., Jin, Y., Unverzagt, F. W., Liang, C., Hall, K. S., Cao, J., and Hendrie, H. C. (2012) 'Selenium level and depressive symptoms in a rural elderly Chinese Cohort', *BMC Psychiatry*, 12(1), pp. 1-8.

- Gao, X., Sanderson, S. M., Dai, Z., Reid, M. A., Cooper, D. E., et al. (2019) 'Dietary methionine influences therapy in mouse cancer models and alters human metabolism', *Nature*, 572(7769), pp. 397-401.
- Gautam, M., Agrawal, M., Gautam, M., Sharma, P., Gautam, A. S. and Gautam, S. (2012) 'Role of antioxidants in generalised anxiety disorder and depression', *Indian Journal of Psychiatry*, 54(3), p. 244.
- Gawlik-Kotelnicka O, Skowrońska A, Margulska A, Czarnecka-Chrebelska KH, Łoniewski I, Skonieczna-Żydecka K, Strzelecki D. (2021) The Influence of Probiotic Supplementation on Depressive Symptoms, Inflammation, and Oxidative Stress Parameters and Fecal Microbiota in Patients with Depression Depending on Metabolic Syndrome Comorbidity-PRO-DEMET Randomized Study Protocol. *J Clin Med*, 10(7):1342. <https://doi.org/10.3390/jcm10071342>
- Ge H, Yang T, Sun J, Zhang D. (2020) Associations between dietary carotenoid intakes and the risk of depressive symptoms. *Food Nutr Res*, 64. <https://doi.org/10.29219/fnr.v64.3920> PMID: 33447180; PMCID: PMC7778430.
- Ghaleiha, A., Davari, H., Jahangard, L., Haghghi, M., Ahmadpanah, M., Seifrabie, M. A., Bajoghli, H., Holsboer-Trachsler, E. and Brand, S. (2016) 'Adjuvant thiamine improved standard treatment in patients with major depressive disorder: results from a randomized, double-blind, and placebo-controlled clinical trial', *European Archives of Psychiatry and Clinical Neuroscience*, 266(8), pp. 695-702.
- Gilbody, S., Lightfoot, T. and Sheldon, T. (2007) 'Is low folate a risk factor for depression? A meta-analysis and exploration of heterogeneity', *Journal of Epidemiology & Community Health*, 61(7), pp. 631-637.
- Gilsing, A. M., Crowe, F. L., Lloyd-Wright, Z., Sanders, T. A., Appleby, P. N., Allen, N. E. and Key, T. J. (2010) 'Serum concentrations of vitamin B12 and folate in British male omnivores, vegetarians and vegans: results from a cross-sectional analysis of the EPIC-Oxford cohort study', *European Journal of Clinical Nutrition*, 64(9), pp. 933-939.
- Gomez-Pinilla, F. and Nguyen, T. T. (2012) 'Natural mood foods: the actions of polyphenols against psychiatric and cognitive disorders', *Nutritional Neuroscience*, 15(3), pp.127-133.
- Grosso, G. (2018) 'Effects of Polyphenol-Rich Foods on Human Health', *Nutrients*, 10(8), p. 1089.
- Grosso, G., Micek, A., Marventano, S., Castellano, S., Mistretta, A., Pajak, A. and Galvano, F. (2016) 'Dietary n-3 PUFA, fish consumption and depression: A systematic review and meta-analysis of observational studies', *Journal of Affective Disorders*, 205, pp. 269-281.
- Grubman, A. and White, A. R. (2014) 'Copper as a key regulator of cell signalling pathways', *Expert Reviews in Molecular Medicine*, 16, p. 11.
- Guo, X., Park, Y., Freedman, N. D., Sinha, R., Hollenbeck, A. R., Blair, A. and Chen, H. (2014) 'Sweetened beverages, coffee, and tea and depression risk among older US adults', *PloS one*, 9(4), 94715.

Halsted JA, Carroll J, Dehghani A, Loghmani M, Prasad AS. (1960) Serum vitamin B12 concentration in dietary deficiency. *Am J Clin Nutr*, 8(3):374-6.

<https://doi.org/10.1093/ajcn/8.3.374>

Hill, C. (2018) 20 Foods That Are High in Vitamin C

<https://www.healthline.com/nutrition/vitamin-c-foods/>

Hills, J. (2019) Top Signs of Iron Deficiency and How to Increase Iron Levels in Your Blood

<https://www.healthyandnaturalworld.com/how-to-increase-iron-levels-in-your-blood/>

Higuera V. (2021) Everything You Need to Know About Depression (Major Depressive Disorder) <https://www.healthline.com/health/depression>

Ho, N. and Prasad, V. (2013) 'Probiotics, prebiotics, synbiotics and naturally fermented foods: why more may be more', *Annals of Gastroenterology*, 26(3), pp. 277–278.

Hopp, C. and Shurtleff, D. (2017) S-Adenosyl-L-Methionine (SAME): In Depth.

<https://www.nccih.nih.gov/health/sadenosylmethionine-same-in-depth/>

Hu, D., Cheng, L. and Jiang, W. (2019) 'Sugar-sweetened beverages consumption and the risk of depression: A meta-analysis of observational studies', *Journal of Affective Disorders*, 245, pp. 348-355.

Huang, Q., Liu, H., Suzuki, K., Ma, S. and Liu, C. (2019) 'Linking What We Eat to Our Mood: A Review of Diet, Dietary Antioxidants, and Depression', *Antioxidants*, 8(9), p. 376.

Huang, S., Zhang, J., Wang, L. and Huang, L. (2013) 'Effect of abiotic stress on the abundance of different vitamin B6 vitamers in tobacco plants', *Plant Physiology and Biochemistry*, 66, pp. 63-67.

Hussein, D.A., Ahmed, G.S., Ahmed, S.F., Salih, R.Q., Kakamad, F.H., et al. (2022). Pattern of vitamin D deficiency in a Middle Eastern population: A cross-sectional study.

International Journal of Functional Nutrition, 3, 7. <https://doi.org/10.3892/ijfn.2022.30>

Hvas, A. M., Juul, S., Bech, P. and Nexø, E. (2004) 'Vitamin B6 level is associated with symptoms of depression', *Psychotherapy and Psychosomatics*, 73(6), pp. 340-343.

Jacka, F. N., O'Neil, A., Opie, R., Itsiopoulos, C., Cotton, S., Mohebbi, M., Castle, D., Dash, S., Mihalopoulos, C., Chatterton, M. L., Brazionis, L., M. Dean, O., M. Hodge, A. and Berk, M. (2017) 'A randomised controlled trial of dietary improvement for adults with major depression (the 'SMILES'trial)', *BMC Medicine*, 15(1), pp. 1-13.

Jelodar, G., Javid, Z., Sahraian, A. and Jelodar, S. (2018) 'Saffron improved depression and reduced homocysteine level in patients with major depression: A Randomized, double-blind study', *Avicenna Journal of Phytomedicine*, 8(1), 43.

Johnson, J. (2019) What foods are good for helping depression

<https://www.medicalnewstoday.com/articles/318428/>

Jones, D. (2019) 7 Healthy Foods That Are High in Vitamin D

<https://www.healthline.com/nutrition/9-foods-high-in-vitamin-d>

Julson, E. (2018) Foods That Are High in Niacin (Vitamin B3)

https://www.healthline.com/nutrition/foods-high-in-niacin#TOC_TITLE_HDR_2/

- Jurowski, K., Szewczyk, B., Nowak, G., and Piekoszewski, W. (2014) 'Biological consequences of zinc deficiency in the pathomechanisms of selected diseases', *Journal of Biological Inorganic Chemistry*, 19(7), pp. 1069-1079.
- Kanekar, S., Bogdanova, O. V., Olson, P. R., Sung, Y. H., D'Anci, K. E. and Renshaw, P. F. (2015) 'Hypobaric hypoxia induces depression-like behavior in female Sprague-Dawley rats, but not in males', *High Altitude Medicine & Biology*, 16(1), pp. 52-60.
- Kang J, Wang D, Duan Y, Zhai L, Shi L, Guo F. (2020) Aerobic Exercise Prevents Depression via Alleviating Hippocampus Injury in Chronic Stressed Depression Rats. *Brain Sci*, 11(1):9. <https://doi.org/10.3390/brainsci11010009>
- Kaviani, M., Nikooyeh, B., Zand, H., Yaghmaei, P. and Neyestani, T. R. (2020) 'Effects of vitamin D supplementation on depression and some involved neurotransmitters', *Journal of Affective Disorders*, 269, pp. 28-35.
- Khan, Sh. (2018) 7 Iron-Rich Foods to Combat Anaemia. www.news18.com/news/indiwo/food-indiwo-7-iron-rich-foods-to-combat-anaemia-1641725.html/
- Kim, N. R., Kim, H. Y., Kim, M. H., Kim, H. M. and Jeong, H. J. (2016) 'Improvement of depressive behavior by Sweetme Sweet Pumpkin™ and its active compound, β -carotene', *Life Sciences*, 147, pp. 39-45.
- Kim, T. H., Choi, J. Y., Lee, H. H., and Park, Y. (2015) 'Associations between dietary pattern and depression in Korean adolescent girls', *Journal of Pediatric and Adolescent Gynecology*, 28(6), pp. 533-537.
- Knüppel, A., Shipley, M. J., Llewellyn, C. H. and Brunner, E. J. (2017) 'Sugar intake from sweet food and beverages, common mental disorder and depression: prospective findings from the Whitehall II study', *Scientific Reports*, 7(1), pp. 1-10.
- Kocot, J., Luchowska-Kocot, D., Kielczykowska, M., Musik, I. and Kurzepa, J. (2017) 'Does vitamin C influence neurodegenerative diseases and psychiatric disorders?', *Nutrients*, 9(7), p. 659.
- Kosheleva, O. V. and Kodentsova, V. M. (2013) 'Vitamin C in fruits and vegetables', *Voprosy Pitaniia*, 82(3), pp. 45-52.
- Krupa, K., Fritz, K. and Parmar, M. (2022) *Omega-3 Fatty Acids*. Treasure Island (FL): StatPearls Publishing. <https://www.ncbi.nlm.nih.gov/books/NBK564314/>
- Lafourcade, M., Larrieu, T., Mato, S., Duffaud, A., Sepers, M., Matias, I., De Smedt-Peyrusse, V., Labrousse, V. F., Bretillon, L., Matute, C., Rodríguez-Puertas, R., Layé, S., and Manzoni, O. J. (2011) 'Nutritional omega-3 deficiency abolishes endocannabinoid-mediated neuronal functions', *Nature Neuroscience*, 14(3), pp. 345-350.
- Lagiou, P., Sandin, S., Lof, M., Trichopoulos, D., Adami, H. O. and Weiderpass, E. (2012) 'Low carbohydrate-high protein diet and incidence of cardiovascular diseases in Swedish women: prospective cohort study', *British Medical Journal*, 344, p. e4026.
- Lande, R. G. (2020) 'Nutraceutical augmentation strategies for depression: a narrative review', *Journal of Osteopathic Medicine*, 120(2), pp. 100-106.

Lange, K., Nakamura, Y., Zhao, H., Bai, D. and Wang, H. (2021) 'Are omega-3 fatty acids efficacious in the treatment of depression? A review', *Journal of Food Bioactives*, 14, pp. 10-19.

Lanuza, F., Petermann-Rocha, F., Celis-Morales, C., Concha-Cisternas, Y., Nazar, G., Troncoso-Pantoja, C., . . . Meroño, T. (2022). A healthy eating score is inversely associated with depression in older adults: Results from the Chilean National Health Survey 2016–2017. *Public Health Nutrition*, 25(10), 2864-2875. doi:10.1017/S1368980021004869

Larsen, T. M., Dalskov, S. M., van Baak, M., Jebb, S. A., Papadaki, A., Pfeiffer, A. F., Martinez, J. A., Handjieva-Darlenska, T., Kunešová, M., Pihlsgård, M., Stender, S., Holst, C., Saris, W. H. M. and Astrup, A. (2010) 'Diets with high or low protein content and glycemic index for weight-loss maintenance', *New England Journal of Medicine*, 363(22), pp. 2102-2113.

Lassale, C., Batty, G. D., Baghdadli, A., Jacka, F., Sánchez-Villegas, A., Kivimäki, M. and Akbaraly, T. (2019) 'Healthy dietary indices and risk of depressive outcomes: a systematic review and meta-analysis of observational studies', *Molecular Psychiatry*, 24(7), pp. 965-986.

Leary, P. F., Zamfirova, I., Au, J. and McCracken, W. H. (2017) 'Effect of latitude on vitamin D levels', *Journal of Osteopathic Medicine*, 117(7), pp. 433-439.

Li, D. and Li, Y. (2019) 'Associations of α -carotenoid and β -carotenoid with depressive symptoms in late midlife women', *Journal of Affective Disorders*, 256, pp. 424-430.

Li, F., Hullar, M. A., Schwarz, Y. and Lampe, J. W. (2009) 'Human gut bacterial communities are altered by addition of cruciferous vegetables to a controlled fruit- and vegetable-free diet', *The Journal of Nutrition*, 139(9), pp. 1685–1691.

Li, G., Mbuagbaw, L., Samaan, Z., Falavigna, M., Zhang, S., Adachi, J. D., ... & Thabane, L. (2014). Efficacy of vitamin D supplementation in depression in adults: a systematic review. *The Journal of Clinical Endocrinology & Metabolism*, 99(3), 757-767.

Li, Z., Li, B., Song, X., and Zhang, D. (2017) 'Dietary zinc and iron intake and risk of depression: A meta-analysis', *Psychiatry Research*, 251, pp. 41-47.

Li, Z., Wang, W., Xin, X., Song, X., and Zhang, D. (2018) 'Association of total zinc, iron, copper and selenium intakes with depression in the US adults', *Journal of Affective Disorders*, 228, pp. 68-74.

Liao, Y., Xie, B., Zhang, H., He, Q., Guo, L., Subramaniapillai, M., Fan, B., Lu, C., and McIntyer, R.S. (2019) 'Efficacy of omega-3 PUFAs in depression: A meta-analysis', *Translational Psychiatry*, 9(1), p. 190.

Libuda, L., Antel, J., Hebebrand, J., & Föcker, M. (2017). Nutrition and mental diseases: focus depressive disorders. *Der Nervenarzt*, 88, 87-101.

Lindblad, M., Tveden-Nyborg, P. and Lykkesfeldt, J. (2013) 'Regulation of vitamin C homeostasis during deficiency', *Nutrients*, 5(8), pp. 2860-2879.

Link, R. (2017) Can Vitamin D Help You Lose Weight
<https://www.healthline.com/nutrition/vitamin-d-weight-loss/>

Lipszyc, P. S., Cremaschi, G. A., Zubilete, M. Z., Bertolino, M. L. A., Capani, F., Genaro, A. M. and Wald, M. R. (2013) 'Niacin modulates pro-inflammatory cytokine secretion. A potential mechanism involved in its anti-atherosclerotic effect', *The Open Cardiovascular Medicine Journal*, 7, pp. 90-98.

Liu, T., Zhong, S., Liao, X., Chen, J., He, T., Lai, S. and Jia, Y. (2015) 'A meta-analysis of oxidative stress markers in depression', *PloS one*, 10(10), p. 0138904.

Ljungberg, T., Bondza, E. and Lethin, C. (2020) 'Evidence of the Importance of Dietary Habits Regarding Depressive Symptoms and Depression', *International Journal of Environmental Research and Public Health*, 17(5), p. 1616.

<https://doi.org/10.3390/ijerph17051616>

Lukić, I., Getselter, D., Koren, O. and Elliott, E. (2019) 'Role of tryptophan in microbiota-induced depressive-like behavior: evidence from tryptophan depletion study', *Frontiers in Behavioral Neuroscience*, 13, 123.

Manosso, L. M., Neis, V. B., Moretti, M., Daufenbach, J. F., Freitas, A. E., Colla, A. R. and Rodrigues, A. L. S. (2013) 'Antidepressant-like effect of α -tocopherol in a mouse model of depressive-like behavior induced by TNF- α ', *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 46, pp. 48-57.

Marx W., Hoare E., Jacka F. (2019) *Diet and Depression: From Epidemiology to Novel Therapeutics*, IN: Editor(s): João Quevedo, André F. Carvalho, Carlos A. Zarate, *Neurobiology of Depression*, Academic Press, Pages 285-292. <https://doi.org/10.1016/B978-0-12-813333-0.00025-1>

Marx W, Moseley G, Berk M, Jacka F. (2017) Nutritional psychiatry: the present state of the evidence. *Proc Nutr Soc*, 76(4):427-436. <https://doi.org/10.1017/S0029665117002026>

Maserejian, N. N., Hall, S. A., and McKinlay, J. B. (2012) 'Low dietary or supplemental zinc is associated with depression symptoms among women, but not men, in a population-based epidemiological survey', *Journal of Affective Disorders*, 136(3), pp. 781-788.

Menon, V., Kar, S. K., Suthar, N., & Nebhinani, N. (2020). Vitamin D and depression: a critical appraisal of the evidence and future directions. *Indian Journal of Psychological Medicine*, 42(1), 11-21.

Miki, T., Kochi, T., Eguchi, M., Kuwahara, K., Tsuruoka, H., et al. (2015) 'Dietary intake of minerals in relation to depressive symptoms in Japanese employees: the Furukawa Nutrition and Health Study', *Nutrition*, 31(5), pp. 686-690.

Mikkelsen, K., Stojanovska, L. and Apostolopoulos, V. (2016) 'The effects of vitamin B in depression', *Current Medicinal Chemistry*, 23(38), pp. 4317-4337.

Mikkelsen, K., Stojanovska, L., Prakash, M. and Apostolopoulos, V. (2017) 'The effects of vitamin B on the immune/cytokine network and their involvement in depression', *Maturitas*, 96, pp. 58-71.

Milaneschi, Y., Bandinelli, S., Penninx, B. W., Corsi, A. M., Lauretani, F., Vazzana, R., Semba, R. D., Guralnik, J. M. and Ferrucci, L. (2012) 'The relationship between plasma carotenoids and depressive symptoms in older persons', *The World Journal of Biological Psychiatry*, 13(8), pp. 588-598.

- Milani, A., Basirnejad, M., Shahbazi, S., and Bolhassani, A. (2017) 'Carotenoids: biochemistry, pharmacology and treatment', *British Journal of Pharmacology*, 174(11), pp. 1290-1324.
- Miyazawa, T., Burdeos, G. C., Itaya, M., Nakagawa, K. and Miyazawa, T. (2019) 'Vitamin E: regulatory redox interactions', *International Union of Biochemistry and Molecular Biology Life*, 71(4), pp. 430-441.
- Młyniec, K., Gawęł, M., Doboszevska, U., Starowicz, G., Pytka, K., Davies, C. L., and Budziszewska, B. (2015) 'Essential elements in depression and anxiety. Part II', *Pharmacological Reports*, 67(2), pp. 187-194.
- Młyniec, K., Gawęł, M., Librowski, T., Reczyński, W., Bystrowska, B., and Holst, B. (2015) 'Investigation of the GPR39 zinc receptor following inhibition of monoaminergic neurotransmission and potentialization of glutamatergic neurotransmission', *Brain Research Bulletin*, 115, pp. 23-29.
- Moludi J, Moradinazar M, Hamzeh B, Najafi F, Soleimani D, Pasdar Y. (2020) Depression Relationship with Dietary Patterns and Dietary Inflammatory Index in Women: Result from Ravansar Cohort Study. *Neuropsychiatr Dis Treat*, 16:1595-1603.
<https://doi.org/10.2147/NDT.S255912>
- Moradi, Y., Albatineh, A. N., Mahmoodi, H. and Gheshlagh, R. G. (2021) 'The relationship between depression and risk of metabolic syndrome: a meta-analysis of observational studies', *Clinical Diabetes and Endocrinology*, 7(1), pp. 1-12.
- Moritz, B., Schmitz, A. E., Rodrigues, A. L. S., Dafre, A. L. and Cunha, M. P. (2020) 'The role of vitamin C in stress-related disorders', *The Journal of Nutritional Biochemistry*, 85, p. 108459.
- Mozaffari, H., Siassi, F., Guilani, B., Askari, M. and Azadbakht, L. (2020) 'Association of dietary acid-base load and psychological disorders among Iranian women: A cross-sectional study', *Complementary Therapies in Medicine*, 53, 102503.
- Müller, C. P., Reichel, M., Mühle, C., Rhein, C., Gulbins, E. and Kornhuber, J. (2015) 'Brain membrane lipids in major depression and anxiety disorders', *Biochimica et Biophysica Acta (BBA)-Molecular and Cell Biology of Lipids*, 1851(8), pp. 1052-1065.
- Muller, D. P. (2010) 'Vitamin E and neurological function', *Molecular Nutrition & Food Research*, 54(5), pp. 710-718.
- Nauser, T., Koppenol, W. H. and Schöneich, C. (2015) 'Protein thiyl radical reactions and product formation: a kinetic simulation', *Free Radical Biology and Medicine*, 80, pp. 158-163.
- Ni, M., You, Y., Chen, J. and Zhang, L. (2018) 'Copper in depressive disorder: a systematic review and meta-analysis of observational studies', *Psychiatry Research*, 267, pp. 506-515.
- NIH. (2011) Prostate Cancer Risk from Vitamin E Supplements. <https://www.nih.gov/news-events/nih-research-matters/prostate-cancer-risk-vitamin-e-supplements/>
- NIH. (2022) Vitamin D: Fact Sheet for Health Professionals
<https://ods.od.nih.gov/factsheets/VitaminD-HealthProfessional>

Nikseresht, S., Etebary, S., Karimian, M., Nabavizadeh, F., Zarrindast, M. R. and Sadeghipour, H. R. (2012) 'Acute administration of Zn, Mg, and thiamine improves postpartum depression conditions in mice', *Archives of Iranian Medicine*, 15(5), pp. 306-311.

Noorazar, S. G., Ranjbar, F., Nemati, N., Yasamineh, N., and Kalejahi, P. (2015) 'Relationship between severity of depression symptoms and iron deficiency anemia in women with major depressive disorder', *Journal of Research in Clinical Medicine*, 3(4), pp. 219-224.

O'Leary, F. and Samman, S. (2010) 'Vitamin B12 in health and disease', *Nutrients*, 2(3), pp. 299-316.

Oh, J., Yun, K., Chae, J. H. and Kim, T. S. (2020) 'Association Between Macronutrients Intake and Depression in the United States and South Korea', *Frontiers in Psychiatry*, 11, p. 207.

Olsen, N. (2019) 20 Foods Rich in Selenium. <https://www.healthline.com/health/selenium-foods/>

Olza, J., Aranceta-Bartrina, J., González-Gross, M., Ortega, R. M., Serra-Majem, L., Varela-Moreiras, G. and Gil, Á. (2017) 'Reported Dietary Intake and Food Sources of Zinc, Selenium, and Vitamins A, E and C in the Spanish Population: Findings from the ANIBES Study', *Nutrients*, 9(7), p. 697.

Pamuk, G. E., Top, M. Ş., Uyanık, M. Ş., Köker, H., Akker, M., Ak, R., Yürekli, Ö. A. and Çelik, Y. (2016) 'Is iron-deficiency anemia associated with migraine? Is there a role for anxiety and depression?', *Wiener Klinische Wochenschrift*, 128(8), pp. 576-580.

Parletta, N., Zarnowiecki, D., Cho, J., Wilson, A., Procter, N., Gordon, A., Bogomolova, S., O'Dea, K., Strachan, J., Ballestrin, M., Champion, A. and Meyer, B. J. (2016) 'People with schizophrenia and depression have a low omega-3 index', *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 110, pp. 42-47.

Pasco, J. A., Jacka, F. N., Williams, L. J., Evans-Cleverdon, M., et al. (2012) 'Dietary selenium and major depression: a nested case-control study', *Complementary Therapies in Medicine*, 20(3), pp. 119-123.

Pathak, L., Agrawal, Y. and Dhir, A. (2013) 'Natural polyphenols in the management of major depression', *Expert Opinion on Investigational Drugs*, 22(7), pp. 863-880.

Pawlak, R., Lester, S. E. and Babatunde, T. (2014) 'The prevalence of cobalamin deficiency among vegetarians assessed by serum vitamin B12: a review of literature', *European Journal of Clinical Nutrition*, 68(5), pp. 541-548.

Peirce, J. M. and Alviña, K. (2019) 'The role of inflammation and the gut microbiome in depression and anxiety', *Journal of Neuroscience Research*, 97(10), pp. 1223-1241.

Permoda-Osip, A., Dorszewska, J., Skibinska, M., Chlopocka-Wozniak, M. and Rybakowski, J. K. (2013) 'Hyperhomocysteinemia in bipolar depression: clinical and biochemical correlates', *Neuropsychobiology*, 68(4), pp. 193-196.

Petre, A. (2017) 5 Ways Restricting Calories Can Be Harmful. <https://www.healthline.com/nutrition/calorie-restriction-risks/>

Petrović-Oggiano, G., Debeljak-Martačić, J., Ranković, S., Pokimica, B., Mirić, A., Glibetić, M. and Popović, T. (2020) 'The effect of walnut consumption on n-3 fatty acid profile of healthy people living in a non-Mediterranean West Balkan country, a small-scale randomized study', *Nutrients*, 12(1), p. 192.

Pfaender, S., Föhr, K., Lutz, A. K., Putz, S., Achberger, K., Linta, L., Liebau, S., Boeckers, T. M. and Grabrucker, A. M. (2016) 'Cellular zinc homeostasis contributes to neuronal differentiation in human induced pluripotent stem cells', *Neural Plasticity*, 2016.

Pierson, H., Yang, H. and Lutsenko, S. (2019) 'Copper Transport and Disease: What Can We Learn from Organoids?', *Annual Review of Nutrition*, 39, pp. 75–94.

Prakash, R., Gandotra, S., Singh, L. K., Das, B. and Lakra, A. (2008) 'Rapid resolution of delusional parasitosis in pellagra with niacin augmentation therapy', *General Hospital Psychiatry*, 30(6), pp. 581-584.

Pratt E. (2019) Kidney Damage and Other Health Problems That Occur from Too Much Vitamin D. <https://www.healthline.com/health-news/too-much-vitamin-d-can-lead-to-kidney-failure/>

Prohan, M., Amani, R., Nematpour, S., Jomehzadeh, N. and Haghhighizadeh, M. H. (2014) 'Total antioxidant capacity of diet and serum, dietary antioxidant vitamins intake, and serum hs-CRP levels in relation to depression scales in university male students', *Redox Report*, 19(3), pp. 133-139.

Psaltopoulou, T., Sergentanis, T. N., Panagiotakos, D. B., Sergentanis, I. N., Kostis, R. and Scarmeas, N. (2013) 'Mediterranean diet, stroke, cognitive impairment, and depression: a meta-analysis', *Annals of Neurology*, 74(4), pp. 580-591.

Ramirez, J., Guarner, F., Bustos Fernandez, L., Maruy, A., Sdepanian, V. L. and Cohen, H. (2020) 'Antibiotics as Major Disruptors of Gut Microbiota', *Frontiers in Cellular and Infection Microbiology*, 10, p. 572912.

Ramos, P., Santos, A., Pinto, N. R., Mendes, R., Magalhães, T. and Almeida, A. (2014) 'Anatomical region differences and age-related changes in copper, zinc, and manganese levels in the human brain', *Biological Trace Element Research*, 161(2), pp. 190-201.

Razzaque, M. S. (2018) 'Magnesium: Are We Consuming Enough?', *Nutrients*, 10(12), p. 1863.

Reinberg, S. (2013) Too Much Fish Oil Might Boost Prostate Cancer Risk. <https://www.webmd.com/prostate-cancer/news/20130710/too-much-fish-oil-might-boost-prostate-cancer-risk-study-says/>

Reininghaus EZ, Platzer M, Kohlhammer-Dohr A, Hamm C, Mörkl S, Lenger M, Bastiaanssen TFS, Dalkner N. (2020) PROVIT: Supplementary Probiotic Treatment and Vitamin B7 in Depression-A Randomized Controlled Trial. *Nutrients*, 12(11):3422. <https://doi.org/10.3390/nu12113422>

Richardson, A. C., Heath, A. L. M., Haszard, J. J., Polak, M. A., Houghton, L. A., and Conner, T. S. (2015) 'Higher body iron is associated with greater depression symptoms among young adult men but not women: observational data from the daily life study', *Nutrients*, 7(8), pp. 6055-6072.

- Rinninella E, Cintoni M, Raoul P, Lopetuso LR, Scaldaferri F, Pulcini G, Miggiano GAD, Gasbarrini A, Mele MC. (2019) Food Components and Dietary Habits: Keys for a Healthy Gut Microbiota Composition. *Nutrients*, 11(10):2393. <https://doi.org/10.3390/nu11102393>
- Rizvi, S., Raza, S. T., Ahmed, F., Ahmad, A., Abbas, S. and Mahdi, F. (2014) 'The role of vitamin E in human health and some diseases', *Sultan Qaboos University Medical Journal*, 14(2), pp. e157–e165.
- Ronco, A. L., Storz, M. A., Martínez-López, W., Calderón, J. M. and Golomar, W. (2021) 'High dietary acid load is associated with prostate cancer risk: An epidemiological study', *World Cancer Research Journal*, 8, e2119.
- Ross, S. M. (2016) 'Omega-3 fatty acids: clinical applications in the treatment of depression', *Holistic Nursing Practice*, 30(6), pp. 382-385.
- Sánchez-Villegas, A., Álvarez-Pérez, J., Toledo, E., Salas-Salvadó, J., Ortega-Azorín, C., Zomeño, M. D., Vioque, J., Martínez, J. A., et al. (2018) 'Seafood Consumption, Omega-3 Fatty Acids Intake, and Life-Time Prevalence of Depression in the PREDIMED-Plus Trial', *Nutrients*, 10 (12), p. 2000.
- Sanchez-Villegas, A., Zazpe, I., Santiago, S., Perez-Cornago, A., Martinez-Gonzalez, M. A. and Lahortiga-Ramos, F. (2018) 'Added sugars and sugar-sweetened beverage consumption, dietary carbohydrate index and depression risk in the Seguimiento Universidad de Navarra (SUN) Project', *British Journal of Nutrition*, 119(2), pp. 211-221.
- Saremi, A. and Arora, R. (2010) 'Vitamin E and cardiovascular disease', *American Journal of Therapeutics*, 17(3), pp. 56-65.
- Sarris, J., Logan, A. C., Akbaraly, T. N., Amminger, G. P., Balanzá-Martínez, et al. (2015) 'Nutritional medicine as mainstream in psychiatry', *The Lancet Psychiatry*, 2(3), pp. 271-274. [https://doi.org/10.1016/S2215-0366\(14\)00051-0](https://doi.org/10.1016/S2215-0366(14)00051-0)
- Sarris, J., Murphy, J., Mischoulon, D., Papakostas, G. I., Fava, M., Berk, M. and Ng, C. H. (2016) 'Adjunctive nutraceuticals for depression: a systematic review and meta-analyses', *American Journal of Psychiatry*, 173(6), pp. 575-587.
- Sarris, J., O'Neil, A., Coulson, C. E., Schweitzer, I. and Berk, M. (2014) 'Lifestyle medicine for depression', *BMC Psychiatry*, 14(1), pp. 1-13.
- Satała, G., Duszyńska, B., Stachowicz, K., Rafalo, A., Pochwat, B., Luckhart, C., et al. (2016) 'Concentration-dependent dual mode of Zn action at serotonin 5-HT1A receptors: in vitro and in vivo studies', *Molecular Neurobiology*, 53(10), pp. 6869-6881.
- Scheiber, I. F., Mercer, J. F. and Dringen, R. (2014) 'Metabolism and functions of copper in brain', *Progress in Neurobiology*, 116, pp. 33-57.
- Schmid A, Walther B. (2013) Natural vitamin D content in animal products. *Adv Nutr*. 4(4):453-62. <https://doi.org/10.3945/an.113.003780>
- Schwingshackl, L., Morze, J. and Hoffmann, G. (2020) 'Mediterranean diet and health status: Active ingredients and pharmacological mechanisms', *British Journal of Pharmacology*, 177(6), pp. 1241-1257.

- Seppälä, J., Koponen, H., Kautiainen, H., Eriksson, J. G., Kampman, O., Leiviskä, J., et al. (2013) 'Association between vitamin B 12 levels and melancholic depressive symptoms: A Finnish population-based study', *BMC Psychiatry*, 13(1), pp. 1-8.
- Shabbir, F., Patel, A., Mattison, C., Bose, S., Krishnamohan, R., Sweeney, E., Sandhu, S., Nel, W., Rais, A., Sandhu, R., Ngu, N., and Sharma, S. (2013) 'Effect of diet on serotonergic neurotransmission in depression', *Neurochemistry International*, 62(3), pp. 324-329.
- Shen YC, Chang CE, Lin MN, Lin CL. (2021) Vegetarian Diet Is Associated with Lower Risk of Depression in Taiwan. *Nutrients*, 13(4):1059. <https://doi.org/10.3390/nu13041059>
- Sheng, J., Liu, S., Wang, Y., Cui, R. and Zhang, X. (2017) 'The Link between Depression and Chronic Pain: Neural Mechanisms in the Brain', *Neural Plasticity*, 2017, 9724371.
- Shiloh, R., Weizman, A., Weizer, N., Dorfman-Etrog, P. and Munitz, H. (2001) 'Antidepressive effect of pyridoxine (vitamin B6) in neuroleptic-treated schizophrenic patients with co-morbid minor depression--preliminary open-label trial', *Harefuah*, 140(5), pp. 369-373.
- Shivavedi, N., Tej, G. N. V. C., Neogi, K. and Nayak, P. K. (2019) 'Ascorbic acid therapy: a potential strategy against comorbid depression-like behavior in streptozotocin-nicotinamide-induced diabetic rats', *Biomedicine & Pharmacotherapy*, 109, pp. 351-359.
- Skolmowska D, Głowska D. (2022) Effectiveness of Dietary Intervention with Iron and Vitamin C Administered Separately in Improving Iron Status in Young Women. *Int J Environ Res Public Health*, 19(19):11877. <https://doi.org/10.3390/ijerph191911877>
- Smesny, S., Baur, K., Rudolph, N., Nenadic, I. and Sauer, H. (2010) 'Alterations of niacin skin sensitivity in recurrent unipolar depressive disorder', *Journal of Affective Disorders*, 124(3), pp. 335-340.
- Sushma, M. (2019) Prescription for plant-based diet overlooks vitamin B12 deficiency. <https://www.downtoearth.org.in/news/food/prescription-for-plant-based-diet-overlooks-vitamin-b12-deficiency-66193/>
- Swardfager, W., Herrmann, N., Mazereeuw, G., Goldberger, K., Harimoto, T., and Lanctôt, K. L. (2013) 'Zinc in depression: a meta-analysis', *Biological Psychiatry*, 74(12), pp. 872-878.
- Szewczyk, B., Kubera, M. and Nowak, G. (2011) 'The role of zinc in neurodegenerative inflammatory pathways in depression', *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 35(3), pp. 693-701.
- Szewczyk, B., Pałucha-Poniewiera, A., Poleszak, E., Pilc, A., and Nowak, G. (2012) 'Investigational NMDA receptor modulators for depression', *Expert Opinion on Investigational Drugs*, 21(1), pp. 91-102.
- Takeda, A. (2000) 'Movement of zinc and its functional significance in the brain', *Brain Research Reviews*, 34(3), pp. 137-148.
- Takeda, A., Tamano, H., Ogawa, T., Takada, S., Ando, M., Oku, N., and Watanabe, M. (2012) 'Significance of serum glucocorticoid and chelatable zinc in depression and cognition in zinc deficiency', *Behavioural Brain Research*, 226(1), pp. 259-264.

Tan, B. L. and Norhaizan, M. E. (2019) 'Carotenoids: How Effective Are They to Prevent Age-Related Diseases?' *Molecules*, 24(9), p. 1801.

Tello, M. (2020) Diet and depression. <https://www.health.harvard.edu/blog/diet-and-depression-2018022213309/>

Terry, N. and Margolis, K. G. (2016) 'Serotonergic mechanisms regulating the GI tract: experimental evidence and therapeutic relevance', *Gastrointestinal Pharmacology*, pp. 319-342.

Thomas, L. D., Elinder, C. G., Tiselius, H. G., Wolk, A. and Åkesson, A. (2013) 'Ascorbic acid supplements and kidney stone incidence among men: a prospective study', *Journal of the American Medical Association Internal Medicine*, 173(5), pp. 386-388.

Tinsley, G. (2018) Methionine: Functions, Food Sources and Side Effects. <https://www.healthline.com/nutrition/methionine/>

Toxqui, L., De Piero, A., Courtois, V., Bastida, S., Sanchez-Muniz, F. J., and Vaquero, M. P. (2010) 'Iron deficiency and overload. Implications in oxidative stress and cardiovascular health', *Nutricion Hospitalaria*, 25(3), pp. 350-365.

Tveden-Nyborg, P. and Lykkesfeldt, J. (2013) 'Does vitamin C deficiency increase lifestyle-associated vascular disease progression? Evidence based on experimental and clinical studies', *Antioxidants & Redox Signaling*, 19(17), pp. 2084-2104.

Vogiatzoglou, A., Refsum, H., Johnston, C., Smith, S. M., Bradley, K. M., De Jager, C., Budge, M. M. and Smith, A. D. (2008) 'Vitamin B12 status and rate of brain volume loss in community-dwelling elderly', *Neurology*, 71(11), pp. 826-832.

Wallace, C. J. and Milev, R. V. (2021) 'The efficacy, safety, and tolerability of probiotics on depression: Clinical results from an open-label pilot study', *Frontiers in Psychiatry*, 12, 132.

Wang, J., Um, P., Dickerman, B. A. and Liu, J. (2018) 'Zinc, magnesium, selenium and depression: a review of the evidence, Potential Mechanisms and Implications', *Nutrients*, 10(5), p. 584.

Wardhana, Surachmanto, E. S. and Datau, E. A. (2011) 'The role of omega-3 fatty acids contained in olive oil on chronic inflammation', *Acta Medica Indonesiana*, 43(2), pp. 138-143.

Ware M. (2022) Health benefits and nutritional value of spinach. <https://www.medicalnewstoday.com/articles/270609>

Watanabe, F. and Bito, T. (2018) 'Vitamin B12 sources and microbial interaction', *Experimental Biology and Medicine*, 243(2), pp. 148-158.

Watson, S. (2022) Folic Acid: Everything You Need to Know. <https://www.healthline.com/nutrition/folic-acid/>

West, H. (2018) The 10 Best Foods That Are High in Zinc. <https://www.healthline.com/nutrition/best-foods-high-in-zinc/>

Whelan, C. (2017) Vitamin B-6 Rich Foods. <https://www.healthline.com/health/vitamin-b6-foods/>

Woznicki, K. (2010) More Chocolate Means More Depression, or Vice Versa.
<https://www.webmd.com/depression/news/20100426/more-chocolate-means-more-depression-or-vice-versa/>

Wu, S., Bhat, Z. F., Gounder, R. S., Mohamed Ahmed, I. A., Al-Juhaimi, F. Y., Ding, Y. and Bekhit, A. E. D. (2022) 'Effect of Dietary Protein and Processing on Gut Microbiota—A Systematic Review', *Nutrients*, 14(3), 453.